

Rice Lake, September, 2023

Fish Survey of Rice Lake (ID #27-0116), Hennepin County, Minnesota in 2023

Survey Dates: September 18-19, 2023

MnDNR Permit Number: 35657

Submitted to:Rice Lake Area
Association and the
City of Maple Grove



Steve McComas Jo Stuckert Blue Water Science St. Paul, MN 55116

Prepared by:

January 29, 2024

Fish Survey of Rice Lake (ID #27-0116), Hennepin County, Minnesota in 2023

Summary

Rice Lake is an 314 acre lake located in Hennepin County, Minnesota.

On September 18-19, 2023, a fish survey using standard trapnets was conducted for Rice Lake. The objective of the fish survey was to fulfill a Minnesota Department of Natural Resources requirement to conduct a fish survey following a lake drawdown. Added benefits to a fish survey would allow the characterization of the existing fish conditions and to determine if fish densities were high enough to be contributing to the observed poor water quality in Rice Lake.

Results of the 2023 fish survey are shown in Table 1. The effects of a 2023 fish kill in Elm Creek did not appear to significantly reduce the fish population in Rice Lake although bass and carp catch rates were lower compared to 2022. The fish catch was dominated by 6 inch and 10 inch black crappies at extremely high catch rates. A total of twelve fish species were observed.

Table 1. Rice Lake trapnet results for fish surveys conducted in 2008, 2014, 2022, 2023 by Blue Water Science and in 1979, 1985, 1994 by the MnDNR. Fish data are shown as fish/trapnet. YOY = young of the year.

	1979 July 12 (5 nets) (MnDNR)	1985 July 9 (8 nets) (MnDNR)	1994 July 11 (9 nets) (MnDNR)	2008 Aug 19-21 (12 nets) (BWS)	2014 July 17-18 (12 nets) (BWS)	2022 Sept 7-8 (12 nets) (BWS)	2023 Sept 19 (6 nets) (BWS)	DNR Range
Bullhead - Black	99	116	1.7	177	7.6	49	79	0.7 - 26
Bullhead - Brown			0.1					1.4 - 6.6
Bullhead - Yellow	0.4	2.8	0.4	0.9	1.0	5.3	4.8	0.8 - 6.2
Carp		2.1	0.3	3.8	0.3	125	12	1.0 - 3.6
Crappie - Black	1.4	17	35	46	2.8	58	522	1.8 - 21
Crappie - White			0.4					2.5 - 11.6
Dogfish (Bowfin)		0.3	1.1	0.2			0.8	0.5 - 1.7
Golden Shiner				1.8	16	0.1	0.2	NA
Largemouth Bass		0.1	0.4	1.8	0.5	3.8	0.2	0.3 - 1.2
Northern Pike	0.4	3.1	0.3	0.1	1.3	0.5	1.3	NA
Sunfish - Bluegill	2.6	42	40	37	62	7.8	13	7.5 - 63
Sunfish - Green		0.4	0.6		0.1			0.2 - 2.0
Sunfish - Hybrid		9.3	1.9			0.5		NA
Sunfish - Orangespot			0.1					NA
Sunfish - Pumpkinseed		2.1	10	1.0	0.1	0.1	0.5	0.8 - 8.4
Tadpole madtom (small bullhead)	0.2							NA
Walleye						0.1		NA
White Sucker		2	0.2	6.5	0.1	0.2	2.5	0.3 - 2.2
Yellow Perch	3.6	21	8.9	4.2	0.7	8.1	9.5	0.5 - 3.4
Bullhead - Black YOY				16	654	94		NA
Carp YOY					405	4.2		NA
Crappie - Black YOY					1.3	206	1.7	NA
Largemouth bass YOY						0.6		NA
Sunfish - Bluegill YOY					2.7		1.0	NA
Yellow perch YOY						0.4		NA

Fish Survey Summaries from 2008-2023

Objectives of this fish survey were to evaluate the effects of previous winterkills and to evaluate the fish community structure after a winter drawdown. Listed below are observations from previous fish surveys from 2008 through 2023.


- The 2008 fish survey found elevated numbers of adult bullheads and carp and poor water quality (Table 1 and Figure 1). Few adult predator fish such as bass, northern pike, or walleyes were found in 2008.
- In the 2014 fish survey, adult bullheads and carp catch rates were lower compared to the 2008 fish survey (Table 1). Water quality was better in 2014 compared to 2008. The winterkill over the 2013-2014 winter likely reduced the adult population of bullheads and carp. However, young of the year carp and bullheads were abundant in 2014.
- The 2022 fish survey was conducted after a significant winter fishkill over the 2021-2022 winter. Small 6 to 8 inch carp were abundant but so were young-of-the-year black crappies. Also, there was an above average adult largemouth bass population. The 2021-2022 fishkill was likely a partial fishkill. Some fish survived and some fish, like carp, migrated into Rice Lake during the summer.
- The fish survey in 2023 was dominated by black crappies at 522 fish/net (Table 1). Carp
 numbers were above the MnDNR range for a shallow lake like Rice, but were down from
 the carp density of 2022. Young of the year fish numbers were low. It's possible that
 predation from the high density crappie population kept the numbers of small fish very low
 (Table 1).
- Because of the possible recruitment of bullheads and carp from Elm Creek, the chances
 of a long-term "balanced" fish community remain a challenge. The best long term strategy
 is to lower phosphorus concentrations in Elm Creek which would reduce phosphorus
 loading to Rice Lake and improve water quality. After water quality is improved, then fish
 stocking would be more effective and possibly sustaining.





The dominant fish in the Rice Lake 2023 fish survey were black crappies at 522 fish/net (left). Carp were common at 12 fish/net (right).

Rice Lake Water Quality






Figure 1. Seasonal averages (May - September) for Secchi disc transparency (top) and total phosphorus (bottom). Recent fish surveys have been conducted in 2008, 2014, 2022, and 2023. The black line indicates the threshold for impaired or unimpaired water quality. For Secchi disc transparency, values greater than 3.3 feet are unimpaired. For phosphorus, concentrations less than 60 ppb are unimpaired.

Fish Survey of Rice Lake (ID #27-0116), Hennepin County, Minnesota in 2023


Introduction

Rice Lake is a 314-acre lake, located in Hennepin County, Minnesota.

In September of 2023, the Rice Lake Area Association contracted for a fish survey with Blue Water Science with a permit granted from the MnDNR. The objective of the fish survey was to fulfill a Minnesota Department of Natural Resources requirement to conduct a fish survey following a lake drawdown. Added benefits to a fish survey would allow the characterization of the existing fish conditions and to determine if fish densities were high enough to be contributing to the observed poor water quality in Rice Lake.

Methods

Six trapnets were used for a single night to survey fish in Rice Lake. Four MnDNR-style trapnets with a 3 x 5 feet square frame with five hoops, two funnel mouth openings and a 50-feet lead. Net mesh size was 3/8 inch (bar length). Also, two MnDNR-style trapnets with a 4 x 6 feet square frame with five hoops, two funnel mouth openings and a 50-feet lead. Net mesh size was



½ inch (bar length). The trapnets were set on Monday morning September 18, 2023. The nets were fished for the following day and removed (September 19). Trapnet locations are shown in Figure 2 and pictures of a typical trapnet are shown in Figure 3.

Figure 2. Map of standard trapnet sets (red) for 2023.

Trapnetting Method



Figure 3. [top] A trapnet is a live fish trap. Fish run into the 50-foot lead net and follow it back through a series of hoops with funnel mouths. Fish end up in the back hoop.
[middle] A handheld net is used to remove the fish from the back of the trapnet.
[bottom] Fish are transferred to tubs, then they are counted, measured, and released.

Results

Standard Trapnets: A total of 12 fish species were sampled in Rice Lake on September 19, 2023. The fish catch was dominated by extremely high densities of black crappies (Table 2).

Although a fishkill was reported in summer in Elm Creek it was likely a partial fishkill. Adult fish for most of the fish species were observed (Tables 2 and 3). Also, successful spawning occurred in 2023.

Table 2. Total fish sampled for the Rice Lake fish survey conducted on September 19, 2023.

September 2023								
•	Net 1	Net 2	Net 3	Net 4	Net 5	Net 6	Total	Fish/net
Black bullhead							P.	
YOY (4-6 inch)	5	10	1	4	93	2	115	19
Adults		100	96	18	140	6	360	60
Bluegill sunfish								
YOY				6			6	1
Adults	1			8	6	8	23	3.8
Black crappie								
YOY (less than 3 in)					10		10	2
Adults	15	1098	340	149	1498	24	3124	521
Carp								
Adults	12	15	22	16	7	2	74	12.3
Largemouth bass								
Adults	1						1	0.2
Northern pike								
Adults		2		2	1	3	8	1.3
Yellow perch								
YOY (less than 4 in)				25			25	4.2
Adults	1		5	3	15	8	32	5.3
Dogfish			1	1	3		5	0.8
Golden shiner					1		1	0.2
Pumpkinseed sunfish	1		1			1	3	0.5
White sucker		2	1	3	8	1	15	2.5
Yellow bullhead	3	12	3	3	5	3	29	4.8
Total fish caught	39	1,239	470	238	1,787	58	3,831	639
Fish per net	7	207	78	40	298	10	639	106

Fish Size Distribution: Length frequencies for Rice Lake fish species surveyed in 2023 are shown in Table 3. The fish length distribution indicates for most fish species there are several year classes present. This indicates that black crappies, black bullheads, and bluegill sunfish may be surviving the winter conditions or there is significant immigration from outside Rice Lake..

Table 3. Length frequency of fish species (as total length) for the Rice Lake fish survey from the standard trapnets for 2023.

Length (inches)	Black bullhead	Black crappie	Bluegill	Carp	Dogfish	Golden shiner	Largemouth Bass	Northern pike	Pumpkinseed	White sucker	Yellow bullhead	Yellow perch
<3		10	6									
3		2	9									1
3.5		19										11
4	4	17										15
4.5		7										1
5	3	16	1								1	11
5.5	6	91	11									3
6	102	90	5			1						6
6.5	8	3	8						1			1
7	5	1	10									1
7.5	1		14								3	
8		2	13								1	
8.5		2	2									2
9	4	15									3	4
9.5	17	22										1
10	26	29									4	
10.5	8	16									2	
11		6		1						1	1	
11.5		1										
12		1								1	6	
12.5											2	
13											5	
13.5											1	
14				7								
14.5				9								
15				17								
15.5				11								
16				12			1					
16.5				1								
17				10	2					1		
17.5												
18				2	1					5		
18.5										1		
19												
19.5										1		
20										4		
21					1					1		
22					1			1				
23												
24				1								
25												
26				1			1	1				
27							1					
28				1				1				
29								1				
30							1					
31				1				2				
32							1	0				
33	404	252	70	74		_		2	_	4-	00	
Measured	184	350	79	74	5	1	1	8	3	15	29 0	57
Counted	291	2784	0	0	0	0	0	0	0			0
TOTAL	475	3134	79	74	5	1	1	8	3	15	29	57
fish/ net (6 nets)	79.2	522.3	13.2	12.3	0.8	0.2	0.2	1.3	0.5	2.5	4.8	9.5

Turtles: Both painted turtles and snapping turtles were common in Rice Lake.

Table 4. Rice Lake painted turtle and snapping turtle catch per net for 2023.

Trapnets

Партосо	TO ⁻	TAL
Net	Painted	Snapping
1		
2	26	
3	2	
4	3	
5	8	
6	2	1
Total Turtles Caught	41	1
Average number/lift	6.8	0.2



Figure 4. Snapping turtle collected in September 2023.

Interesting Fish Related Observations in 2023





Crappies

Northern pike





Bluegills

Largemouth bass





Dogfish

Carp

Fish Conditions from Surveys in 2008, 2014, and 2022



2008



2014



2022

Fish Survey Summaries from 2008-2023

Objectives of this fish survey were to evaluate the effects of previous winterkills and to evaluate the fish community structure after a winter drawdown. Listed below are observations from previous fish surveys from 2008 through 2023.

- The 2008 fish survey found elevated numbers of adult bullheads and carp and poor water quality (Table 1 and Figure 1). Few adult predator fish such as bass, northern pike, or walleyes were found in 2008.
- In the 2014 fish survey, adult bullheads and carp catch rates were lower compared to the 2008 fish survey (Table 1). Water quality was better in 2014 compared to 2008. The winterkill over the 2013-2014 winter likely reduced the adult population of bullheads and carp. However, young of the year carp and bullheads were abundant in 2014.
- The 2022 fish survey was conducted after a significant winter fishkill over the 2021-2022 winter.
 Small 6 to 8 inch carp were abundant but so were young-of-the-year black crappies. Also, there was an above average adult largemouth bass population. The 2021-2022 fishkill was likely a partial fishkill. Some fish survived and some fish, like carp, migrated into Rice Lake during the summer.
- The fish survey in 2023 was dominated by black crappies at 522 fish/net (Table 1). Carp numbers were above the MnDNR range for a shallow lake like Rice, but were down from the carp density of 2022. Young of the year fish numbers were low. It's possible that predation from the high density crappie population kept the numbers of small fish very low (Table 1).
- Because of the possible recruitment of bullheads and carp from Elm Creek, the chances of
 a long-term "balanced" fish community remain a challenge. The best long term strategy is
 to lower phosphorus concentrations in Elm Creek which would reduce phosphorus loading
 to Rice Lake and improve water quality. After water quality is improved, then fish stocking
 would be more effective and possibly sustaining.

Appendix

Appendix: E-Mail Notification of Fish Survey to Be Conducted

From: mccomas@pclink.com < mccomas@pclink.com >

Sent: Friday, September 15, 2023 11:21 AM

To: Daryl Ellison <daryl.ellison@state.mn.us>; Capt. Jason Peterson <jason.r.peterson@state.mn.us>

Cc: George Schneider <dukewolves@comcast.net>

Subject: Fish Survey in Rice Lake (27-011600), Hennepin Co September 18-19, 2023

Hello all,

Blue Water Science will be conducting a fish survey in Rice Lake (MN ID 27-011600), Hennepin County, starting on Monday, September 18, 2023. We will set 6 trap nets in the lake. The nets will be monitored daily and removed on Wednesday, September 19, 2023 and all fish will be weighed, measured, and returned to the lake. The nets will be removed from the lake on Wednesday, September 19, 2023. The fish survey is sponsored by the Rice Lake Association with the objectives of characterizing the existing fish community structure and assessing potential impacts of fish on water quality.

This survey is being conducted under the permit number: 35657.

Thank you,
Steve McComas
BLUE WATER SCIENCE
550 South Snelling Avenue
St. Paul, MN 55116

651 231 2778 - cell mccomas@pclink.com

Appendix: 2022 Rice Lake Trapnet Results

Total fish sampled for the Rice Lake fish survey conducted on September 6-8, 2022.

September 2022														
	Ne	et 1	Ne	et 2	Ne	t 3	Ne	et 4	Ne	t 5	Ne	et 6	Total	Fish/
	Day 1	Day 2		net										
Black bullhead														
YOY (4-6 inch)	500	280	35	38	67	16	33		19		63	80	1131	94
Adults	128	125	15	5	66	83	4	1	70	20	5	70	592	49
Bluegill sunfish														
YOY													0	0
Adults	20	32	3	10	2	8	5		8		4	1	93	7.8
Black crappie			•		•				•					
YOY (less than 3 in)	93	320	440	86	54	520	73		1	26	56	420	2,089	174
Adults	39	68	9	3	7	8	3		94	410	48	3	692	58
Carp														
YOY		50											50	4.2
2 year old fish	408	290	21	57	170	2	240	181	27	2	43	36	1477	123
Adults	1	1	2	1			1	1	3	5	1	4	20	1.7
Largemouth bass														
YOY (less than 5 in)			1		6								7	0.6
Adults	13	16	3			2		2	3	2	5		46	3.8
Northern pike														
Adults		1				3			1	1			6	0.5
Yellow perch			•		•				•					
YOY (less than 4 in)	1						4						5	0.4
Adults	9	20	1		1	6	12	29	5	7	1	6	97	8.1
Golden shiner							1						1	0.1
Hybrid sunfish	1	2	1		1		1						6	0.5
Pumpkinseed sunfish	1												1	0.1
Walleye			1										1	0.1
White sucker		1							1				2	0.2
Yellow bullhead	2	29	4		2		1	1	1	4		19	63	5.3
Total fish caught	1,216	1,235	536	200	376	648	378	215	233	477	226	639	6,379	532
Fish per net	203	206	89	33	63	108	63	36	39	80	38	107	1,065	89

2022: Length frequency of fish species (as total length) for the Rice Lake fish survey from the standard trapnets for 2022.

Rice	Black bullhead	Black crappie	Bluegill		Golden shiner	Hybrid sunfish	LM Bass	Northern pike	Pumpkin- seed	Walleye	White sucker	Yellow bullhead	Yellow perch
<3	1131	2472		71									5
3	1												7
3.5		1			1								5
4	2						3						31
4.5				1			2						3
5				9			3						1
5.5				13			1						1
6			7	20			1						1
6.5		1	20	24		1							1
7	1		47	38		2			1				7
7.5		2	18	25									3
8	4	7	3	17		1						2	7
8.5	8	31	3	6		1							10
9	20	94	1	2		1							12
9.5	47	46		2									6
10	76	27					1					2	2
10.5	28	4					1					3	
11	10	1					3					4	
11.5	1						1				1	2	
12							11					2	
12.5	1						2						
13							3						
13.5							10						
14							5						
14.5							4						
15		1					1						
15.5													
16							1						
16.5													
17													
17.5													
18											1		
18.5													
19													
19.5										1			
20													
20.5													
21								1					
21.5													
22													
22.5								1					
23								1					
23.5													
24				1									
24.5													
25													
26				4									
27				2				1					
28				1									
29				2									
30				5									
31				1				1					
32				1				1					
33				2									
34													
35													
36				1									
Measured		2687	99	248	1	6	53	6	1	1	2	15	102
Counted		94	0	1298	0	0	0	0	0	0	0	0	0
TOTAL	1723	2781	99	1546	1	6	53	6	1	1	2	15	102
fish/ net (12 nets)	144	232	8.3	129	0.1	0.5	4.4	0.5	0.1	0.1	0.2	1.3	8.5

Appendix: 2014 Rice Lake Trapnet Results

Rice Lake trapnet results for the fish survey conducted in July 2014.

					Fish C	aptured (July 17-18	3 2014)					Total	2014	Normal
	Ne	t 1	Ne	t 2	Ne	et 3	Ne	t 4	Ne	t 5	Ne	et 6	Catch	Fish per Net	Range (MnDNR)
	Day 1	Day 2	Day 1	Day 2	Day 1	Day 2	Day 1	Day 2	Day 1	Day 2	Day 1	Day 2		(n=12)	(2)
Black bullhead (Ameiurus melas)	15	13	8	9	5		5		11	8	8	9	91	7.6	2.2 - 60.5
Black crappies (Pomoxis nigromaculatus)	2	4	1	1	3	1	6		4	8	1	2	33	2.8	2.4 - 15.1
Bluegill sunfish (Lepomis macrochirus)	68	51	95	56	44	10	75	1	184	71	82	12	749	62	1.9 - 29.5
Carp (Cyprinus carpio)			1		1		1						3	0.3	1.0 - 3.6
Golden shiner (Notemigonus crysoleucas)	18	17	10	1	9	1	7	1	58	30	15	27	194	16	NA
Green Sunfish (Lepomis cyanellus)				1									1	0.1	0.2 - 2.0
Largemouth bass (Micropterus salmoides)					1		2			1	1	1	6	0.5	0.3 - 1.2
Northern Pike (Esox lucius)			3	5							4	4	16	1.3	NA
Pumpkinseed sunfish (Lepomis gibbosus)											1		1	0.1	0.8 - 8.4
White sucker (Catostomus commersonii)											1		1	0.1	0.3 - 2.2
Yellow bullheads (Ameiurus natalis)			5	5						1		1	12	1.0	0.8 - 6.2
Yellow perch (Perca flavescens)			2	2	1		1			1	1		8	0.7	0.5 - 3.4
TOTAL FISH	103	85	125	80	64	12	97	2	257	120	114	56	1,115	92.9	
Turtles - painted			2	1	13	1	5		1	1	11	1	36	1.5	
Turtles - snapping					2			1			2		5	0.9	
Black bullhead YOY (Ameiurus melas)			4,852	190	2,710	95							7,847	654	NA
Black crappies YOY (Pomoxis nigromaculatus)						5		10					15	1.3	NA
Bluegill sunfish YOY (Lepomis macrochirus)						10	16	6					32	2.7	NA
Carp YOY (Cyprinus carpio)	15	12	756	1,036	626	39	100	12	456	939	25	846	4,862	405	NA

2014: Length frequency of fish species (as total length) for the Rice Lake fish survey from the standard trapnets for 2014 (young of the year bullhead lengths are not shown).

size (inches)	Black bullhead	Bluegills	Carp	Crappies	Golden shiner	Green sunfish	Large- mouth	Northern pike	Pumpkin- seeds	White suckers	Yellow bullhead	Yellow perch
(IIICIICS)	Dullileau				Sillilei	Suminism	bass	PIKE	36603	Suckers	Dullileau	percii
<3		32		15								
3		6	4				1					1
3.5			6									
4		3	28		3		2					1
4.5		3	65	1	9	1			1			
5		2	61	1	10							
5.5		3	65	1	6							
6	4	25	13		53			1			2	1
6.5	1	154	2		38							
7	3	144		1	34			1			1	
7.5	1	43		6	1			4			1	2
8	2	1		8				2				3
8.5				12				2			2	
9	7			3				4			1	
9.5	12							1			3	
10	22							1			2	
10.5	26											
11	13											
11.5												
12												
12.5												
13												
13.5												
14												
14.5												
15							1			1		
15.5												
16												
16.5												
17							1					
18												
19							1					
20												
21												
22												
23												
24			1									
25			1									
26			1									

Appendix: 2008 Rice Lake Trapnet Results

Rice Lake trapnet results for the fish survey conducted in August 2008.

Net	Bluegill	Bullhead Black	Bullhead Black YOY	Bullhead Yellow	Black Crappie	Carp	Dogfish	Golden Shiner	Largemouth Bass	Northern Pike	Pumpkin- seed	Sucker	Yellow Perch
Tuesday (8/1	19/08)												
1	37	141		1	30	4			4			3	4
2	64	648	10	2	42	11	1		1	1	1	9	
3	77	112	90	1	65				2		7	21	3
4	2	2		1	7			13					1
subtotal	180	903	100	5	144	15	1	13	7	1	8	33	8
fish/ net	45	226	25	1.3	36	3.8	0.3	3.3	1.8	0.3	2.0	8.3	2.0
Wednesday	(8/20/08)												
1	61	163		3	46	2			1		1	7	25
2	43	100	24		70	12			1		2	2	2
3	51	236	61		63	4			3		1		5
4	3	16			8	1		5					1
subtotal	158	515	65	3	187	19	0	5	5	0	4	9	33
fish/ net	40	129	16	0.8	47	4.8		1.3	1.3		1.0	2.3	8.3
Thursday (8/	/21/08)												
1	9	49		1	64	3		3	5			17	8
2	56	86		1	78	4	1		4			12	
3	42	555	26	1	70	3			1			7	
4	2	16			6	2		1					1
subtotal	109	706	26	3	218	12	1	4	10	0	0	36	9
fish/ net	27	177	6.5	0.8	55	3.0	0.3	1.0	2.5			9.0	2.3
Total Fish (12 nets)	447	2,124	191	11	549*	46	2	22	22	1	12	78	50
Fish/ Trapnet	37	177	16	0.9	46	3.8	0.2	1.8	1.8	0.1	1.0	6.5	4.2
MnDNR Normal Range*	7.5 - 63	0.7 - 26	NA	0.8 - 6.2	1.8 - 21	1.0 - 3.6	0.5 - 1.7	NA	0.3 - 1.2	NA	0.8 - 8.4	0.3 - 2.2	0.5 - 3.4

^{* 27} YOY black crappies were involved in total count

Rice Lake mini-trapnet results for the fish survey conducted in August 2008.

Net	Bluegill	Bluegill YOY	Bullhead Black	Bullhead Black YOY	Bullhead Yellow	Black Crappie	Black Crappie YOY	Carp	Golden Shiner	Large- mouth Bass	Pumpkin- seed	Sucker	Yellow Perch
Tuesday (8,	/19/08)												
1	18		17	42		31	1	2	6	2			9
2	6	1,620	1			2							1
3*	4	30	7		1	13	16	2		1	1		2
subtotal	28	1,650	25	42	1	46	17	4	6	3	1	0	12
fish/ net	9.3	550	8.3	14	0.3	15	5.7	1.3	2.0	1.0	0.3		4.0
Wednesday	y (8/20/08)												
1	15		1	26		13		3	2				2
2	8	1,500	2										
3	9		13	52		19				1			2
subtotal	32	1,500	16	78		32		3	2	1	0	0	4
fish/ net	11	500	5.3	26		11		1.0	0.7	0.3			1.3
Thursday (8	3/21/08)												
1	3			25		11			1				
2*	4	750				6		1	1	4			2
3	3	50	9	2	2	10	16			4		2	10
subtotal	10	800	8	27	2	27	16	1	2	8	0	2	12
fish/ net	3.3	267	3.0	9.0	0.7	9.0	5.3	0.3	0.7	2.7		0.7	4.0
Total Fish (9 nets)	70	3,950	50	147	3	105	33	8	10	12	1	2	28
Fish/ Mini Trapnet	7.8	439	5.6	16	0.3	12	3.7	0.9	1.1	1.3	0.1	0.2	3.1

*Day 1; net 3: crayfish = 1 Day 3; net 2: shiner minnows = 2

2008: Length frequency of fish species (as total length) for the Rice Lake fish survey from the standard trapnets.

Size Range	Bluegill	Black	Black	Carp	Dogfish	Golden	Largemouth	Northern	Pumpkin-	Sucker	Yellow
(in)		Bullhead	Crappie			Shiner	Bass	Pike	seed		Perch
<3.0											
3.0							2				
3.5							2				
4.0	1						2				
4.5											
5.0	14					3	5		7		
5.5	21								1		
6.0	88	1	3			10					6
6.5	60	1				3					2
7.0	55	4	6			4	1				22
7.5	5	4	9								7
8.0	1	22	87			1					9
8.5		2	42								3
9.0			76								
9.5			4								
10.0		2	4								
10.5		1									
11.0											
11.5											
12							1				
13							1			1	
14 15							1				
16							3			1 6	
17							3			24	
18							2			21	
19										22	
20										2	
21				8						1	
22				9	2						
23				7							
24				11							
25				3							
26				2				1			
27											
28											
29											
30				1				1			
31				1							